

J. Michael Conoyer, M.D. Karen E. Boone, M.D. Robert R. MacDonald III, M.D. J. Matthew Conoyer, M.D., F.A.C.S. Benjamin M. Conoyer, M.D. Matthew P. Page, M.D.

4790 Executive Centre Parkway, St. Peters, MO 63376

Ear Surgery Postoperative Instructions

Special Instructions

- If a protective bandage has been applied, you may remove it the day after surgery.
- Sutures or Steri-Strips may be removed in the office.
- Change the cotton ball in the ear canal at least daily and use ear drops if so instructed.
- Keep the ear dry until the follow up visit.
- Hair may be washed, but keep the ear canal dry by using a cotton ball covered in antibiotic ointment or Vaseline[®].
- Do not drive if you are taking narcotic pain medication or medication for dizziness.

Medication

- Tylenol[®], Motrin[®] or prescribed pain medication may be used as needed.
- Use ear drops or apply Bacitracin[®] ointment to the incision behind the ear, if instructed by your surgeon.
- Take antibiotics by mouth if prescribed by your surgeon.

Activity

- Avoid strenuous activity for at least 3 weeks after surgery.
- Avoid flying for at least 3 weeks after surgery. Discuss with your surgeon when it is OK to fly.
- Dry ear precautions until cleared by your surgeon.
- Avoid hard nose blowing for 3 weeks after surgery.

Diet

• No restrictions.

Postoperative Follow-up Appointment

- One to four weeks as instructed by your surgeon.
- Call 636.441.3100 should you need to reschedule or confirm your appointment time.

What to Expect

- Mild to moderate pain with some swelling.
- Drainage that may be bloody or contain small pieces of the absorbable ear canal packing.
- Decreased hearing that will take several weeks to improve.
- Low grade fever will sometimes follow general anesthesia for the first three days.

When to Call

- Call the office at 636.441.3100 during routine hours for questions, concerns or medication refill requests. Please remember that pain medications will only be refilled during regular office hours.
- After-hours and weekend calls should be limited to urgent and emergent issues such as the following:
 - o Excessive bleeding
 - o Adverse reactions to prescribed medications, such as rash or vomiting
 - Fever >101.4° that does not respond to Tylenol[®] or Motrin[®].
- Should you choose or be instructed to go the ER for any reason, we ask that you go to either St. Joseph Hospital-Lake St. Louis or Barnes-Jewish St. Peters Hospital.