

J. Michael Conoyer, M.D. Karen E. Boone, M.D. Robert R. MacDonald III, M.D. J. Matthew Conoyer, M.D., F.A.C.S. Benjamin M. Conoyer, M.D. Matthew P. Page, M.D.

4790 Executive Centre Parkway, St. Peters, MO 63376

Nasal Septoplasty

Postoperative Instructions

Special Instructions

- Start nasal saline (Ocean[®]) the day of surgery. Use 4-5 times daily until your first follow-up visit. This will help prevent crusting inside your nasal splints and will help you breathe better until their removal.
- Do not blow your nose the first week.
- Do not drive while taking narcotic pain medications.

Medication

- Use prescribed narcotic pain medication as directed. You may supplement with Motrin® as needed.
- Antibiotics will be prescribed if necessary.
- If you were previously taking aspirin or other blood-thinning agents, discuss with your surgeon when to restart these medications.

Activity

• Limited activity for one week. Do not lift >30 pounds. You may do some limited aerobic exercise as tolerated.

Diet

• No restrictions.

Postoperative Follow-up Appointment

- Usually one week after surgery, at which time any nasal splints or packing will be removed.
- Call 636.441.3100 should you need to reschedule or confirm your appointment time.

What to Expect

- Minor bloody discharge from the nose is to be expected for up to one week after surgery. Should you experience a
 true nosebleed, lay in a reclined position, instill 2 sprays of Afrin[®] to each nare, and hold pressure for 10 minutes. Call
 the office at 636.441.3100 should it fail to resolve.
- Discomfort will gradually improve over the first week, and will improve when splints are removed.

When to Call

- Call the office at 636.441.3100 during routine hours for questions, concerns or medication refill requests. Please remember that pain medications will only be refilled during regular office hours.
- After-hours and weekend calls should be limited to urgent and emergent issues such as the following:
 - Excessive bleeding
 - Adverse reactions to prescribed medications, such as rash or vomiting
 - Fever >101.4° that does not respond to Tylenol[®] or Motrin[®]
- Should you choose or be instructed to go the ER for any reason, we ask that you go to either St. Joseph Hospital-Lake St. Louis or Barnes-Jewish St. Peters Hospital.